

REDESCRIPTION OF *GYRAULUS ARGAEICUS* (STURANY 1904) WITH THE DESCRIPTION OF TWO NEW GASTROPOD SPECIES FROM TURKEY (MOLLUSCA: GASTROPODA: BITHYNIIDAE, PLANORBIDAE)

PETER GLÖER¹ & DILIAN GEORGIEV²

¹Schulstrasse 3, D-25491 Hetlingen, Germany

²Department of Ecology and Environmental Conservation, University of Plovdiv, Tzar Assen Str. 24, BG-4000 Plovdiv, Bulgaria

Abstract New records of freshwater snails from Turkey are presented. Two species new for science *Bithynia yildirimi* and *Gyraulus nedylalkovi* are described. In addition the junior author collected *Gyraulus argaeicus* from its type locality, so the anatomy of this species is provided for the first time. Furthermore, new record of *Anisus leucostoma*, confirmed by the study on its anatomy, is given.

Key words *Bithynia*, *Gyraulus*, *Gyraulus argaeicus*, *redescription*, *new species*, *Turkey*.

INTRODUCTION

Six species of genus *Bithynia* are known from Turkey (Glöer & Yıldırım, 2006; Yıldırım, 1999): *B. tentaculata* (Linnaeus 1758), *B. leachii* (Sheppard 1824), *B. pseudemmericia* Schütt 1964, *B. phialensis* (Conrad 1852), *B. badiella* (Küster 1852), and *B. pesicci* Glöer & Yıldırım 2006. The closely related genus *Pseudobithynia* is represented in Turkey by *P. pentheri* (Sturany 1904) (Glöer & Yıldırım, 2006).

According to Yıldırım *et al.* (2008) six *Gyraulus* species are known from Turkey: *Gyraulus albus* (O.F. Müller 1774), *G. laevis* (Alder 1838), *G. piscinarum* (Bourguignat 1852), *G. ehrenbergi* (Beck 1837), *G. euphraticus* (Mousson 1874), *G. parvus* (Say 1817), *G. crista* (Linnaeus 1758), and *G. hebraicus* (Bourguignat 1852). In addition, Glöer & Rähle (2009) recently described *Gyraulus pamphylicus* Glöer & Rähle 2009 from Turkey.

From neighbouring countries some more species of genus *Gyraulus* are known, ie. *G. homsensis* (Dautzenberg 1894) from Syria, (Dautzenberg, 1894: 337), *G. bekaensis* Glöer & Bößneck 2007 from Lebanon (Glöer & Bößneck, 2007: 142–144) and *G. huwaizahensis* Glöer & Naser 2007 from Iraq (Glöer & Naser, 2007: 150–151). Bank (2004) listed following species: *G. albus*, *G. laevis*, and *G. crista* from Greece, and Angelov (2000) mentioned *G. albus*, *G. laevis*, *G. crista* and *G. piscinarum* from Bulgaria.

This paper is aimed at redescribing *Gyraulus argaeicus* and describing two new species: *Bithynia yildirimi* n. sp. and *Gyraulus nedylalkovi* n. sp.

MATERIAL AND METHODS

The snails were collected with a sieve from the banks of the relevant waters. Sampling sites are given in Fig. 1.

The samples were put into ethanol (75%). The dissections and measurements of the genital organs and the shells were carried out using a stereo microscope (Zeiss, Germany). The



Figure 1 The map of study area with marked sampling sites. Sampling sites: 1 *Bithynia yildirimi* n. sp., *Gyraulus nedylalkovi* n. sp., *Anisus leucostoma*; 2 *Gyraulus argaeicus*.

photographs were made with a digital camera system (Leica R8).

The type material will be deposited in the Zoological Museum Hamburg (ZMH), Germany.

RESULTS

Conchological and anatomical investigations revealed two new species which belong to the genera *Bithynia* and *Gyraulus*, respectively.

Family Bithyniidae Troschel 1857

Genus *Bithynia* Leach 1818

Type species: *Bithynia tentaculata* (Linnaeus 1758)

Bithynia yildirimii n. sp.

Material examined 23 exx. from type locality.

Holotype Shell height 4.8 mm, width 2.9 mm, Zoological Museum Hamburg ZMH 79174.

Paratypes 5 exx., ZMH 79175.

Locus typicus Turkey, Mediterranean Sea coast, a swamp east of Kazanli village, N36°48'24.7" E34°47'34.8", 13.08.2009 D. Georgiev leg.

Habitat A swamp near the Mediterranean coast, densely occupied by water and bank vegetation

as *Phragmites australis*, and various bush and trees. Some parts of the swamp completely drying during summer.

Etymology Named after Prof. Dr M.Z. Yıldırım (Eğridir, Isparta), the outstanding expert on freshwater molluscs of Turkey.

Diagnosis The yellowish shell is slim, of 4–5 convex whorls with clear sutures (Fig. 2a). Aperture, also the operculum (Fig. 2b), shows an obtuse angle. Umbilicus slit-like to closed. Nucleus of operculum eccentric. Clear sexual dimorphism shown (Fig. 2a,c). Male shells 4.7–5.1 mm high, 2.9–3.0 mm wide; female shells larger, 5.1–5.4 mm high, 3.6–3.7 mm wide.

Anatomy Penial appendix branches off from distal third of penis and is twice longer than distal part of the penis (Fig. 2d). Flagellum short (Fig. 2e).

Remarks We do not believe that *Bithynia leachii* occurs in Turkey (Yıldırım *et al.*, 2006). This species is distributed in the lowlands of western Europe towards Russia, and the southernmost records known are from Hungary (Glöer & Fehér, 2004). Thus we have to compare *Bithynia yildirimii* sp. nov. with three other *Bithynia* species known from Turkey, i.e. *B. pesicci*, *B. phialensis*, and *B. badiella*. The suture in *B. pesicci*




Figure 2 *Bithynia yildirimii* n. sp.: a holotype, male, shell; b operculum; c paratype, female, shell; d penis in situ; e Penis with flagellum. Parts: fl flagellum; p penis; pa penial appendix.

is very deep and the whorls are swollen, easily distinguishing the latter species from *B. yildirimii*. Furthermore, the flagellum of *B. pesicci* is much shorter than in *B. yildirimii*. The shell of *B. badiella* is spherical, distinct from *B. yildirimii*. The suture of *B. phialensis* is more flattened and the shell is broader than in *B. yildirimii*. It is worth mentioning that the presence of both the aforementioned species, i.e. *B. phialensis* (type locality in Palestine) and *B. badiella* (type locality: Beirut in Turkey), are doubtful.

Genus Gyraulus Charpentier 1837

Type species: *Planorbis albus* O.F. Müller 1774

***Gyraulus argaeicus* (Sturany 1904)**

Material examined 11 exx. from type locality.

Locus typicus Turkey, Lake of Soysali village, 12.08.2009, D. Georgiev leg.

Habitat The Soysali Lake is situated on volcanic rocks, and fed by a big spring with the waters emerging on its northern bank. The water is clear and the lake bottom is covered mainly by gravel with very few sandy or muddy zones. Pollution from garbage from the village of Soysali and cattle watering was observed. The surrounding habitat is typical of steppe in the Anatolian Plateau.

Description The light-corneous shell nearly dull and transparent with fine growth lines. The shell consists of 3–4 whorls, which regularly and rapidly increase with a clearly visible to deep suture. The last whorl is slightly deflected (Fig. 3a). The first whorls are immersed on the underside, forming a deep umbilicus. The shell is 6.1–7.1 mm in diameter and 1.5–1.8 mm in height.

Animal The animal is light grey with a diffuse mantle pigmentation. The prostate gland bears




Figure 3 *Gyraulus argaeicus* (Topotype): **a** the shell; **b** sex tract; **pr** prostate gland; **pht** phallotheca; **prp** praeputium; **vd** vas deferens.

1. *Planorbis (Gyraulus) argaeicus* n. sp. — Schale flach, oben und unten etwas konkav, nahezu glanzlos, von grünlich-gelber Farbe, mit vier rasch anwachsenden, durch eine seichte Naht getrennten Windungen, mit zarten und dicht aneinander gerückten Anwachsstreifen und feinsten Spirallinien; letzter Umgang zuweilen in der Mitte gekielt; Mündung schief ohrförmig, Oberrand stark vorgezogen und mit dem genäherten Spindelrande durch einen Callus verbunden.

Schalbreite 7, Schalenhöhe 2.2 mm; Mündung 3 mm breit und 2.7 mm hoch.

Fundort: Soisaly im Erdschiasgebiete, Kleinasiens (leg. Penther).

Figure 4 Facsimile of the original description (Sturany, 1904: 115–116).

20–22 long diverticula. The phallotheca is nearly twice as long as the praeputium (Fig. 3b).

Remarks *Gyraulus argaeicus* (Sturany 1904), described from Turkey (Soisaly), has not been mentioned since its original description (Sturany,

1904: 115). It was not possible to compare our material with the type material, because the Vienna Museum does not lend holotypes or paratypes for scientific studies. However, the original description (Fig. 4) corresponds well with our specimens.

***Gyraulus nedyalkovi* n. sp.**

Fig. 5a–d

Material examined 14 exx. from type locality.

Holotype Shell height 1.4 mm, width 4.4 mm, Zoological Museum Hamburg ZMH 9176.

Paratypes 5 exx., ZMH 79177.

Locus typicus Turkey, Mediterranean Sea coast, a swamp east of Kazanli village, N36 48 24.7 E34 47 34.8, 13.08.2009 D. Georgiev leg.




Figure 5 *Gyraulus nedyalkovi* n. sp.: **a** holotype (photographed in ethanol), shell; **b** paratype, shell; **c** male copulatory organ; **d** bursa copulatrix and prostate gland. Parts: **bc** bursa copulatrix; **m** muscle; **pr** prostate gland; **pht** phallotheca; **prp** praeputium; **vd** vas deferens.



Figure 6 The shells of *Gyraulus* species from the Near East region: 1 *G. pamphylicus*; 2 *G. hebraicus*; 3 *G. euphraticus*; 4 *G. argaeicus* (topotype); 5 *G. homsensis*; 6 *G. ehrenbergi*; 7 *G. huwaizahensis*; 8 *G. piscinarum*; 9 *G. bekaensis*; 10 *G. nedyalkovi* n. sp.

Table 1 Distinguishing characteristics of *Gyraulus* spp. from the Near East region.

Taxon, type country	max. diameter D [mm]	height of last whorl h [mm]	ratio D: h	keel	no. of whorls	last whorl prominent	umbilicus	no. of prostate diverticules
<i>G. nedyalkovi</i> n. sp. Turkey	3.9	1.0	3.9	no	3–4	yes	wide	18–22
<i>G. pamphylicus</i> , Turkey	7.0	1.0–1.2	7.0–5.8	yes	4	yes	narrow	16–18
<i>G. argaeicus</i> , Turkey	7.0	2.2	3.2	slight	4	yes	wide	18–22
<i>G. hebraicus</i> , Syria	5.0	1.2	4.2	no	4	no	narrow	11–15
<i>G. homsensis</i> , Syria	5.0	1.0	5.0	slight	4.5	yes	wide	20–22
<i>G. piscinarum</i> , Lebanon	4.4	1.1–1.2	4.0–3.7	no	3.5	yes	wide	12–16
<i>G. bekaensis</i> , Lebanon	5.7	1.4–1.5	4.1–3.8	yes	3.5	yes	narrow	18
<i>G. ehrenbergi</i> , Egypt	4.5	1.0	4.5	no	3.5	yes	narrow	14–19
<i>G. huwaizahensis</i> , Iraq	3.0–3.5	1.0	3.0–3.5	no	3.75	yes	narrow	9
<i>G. euphraticus</i> , Iraq	7.0	1.0	7.0	slight	4.5	no	narrow	9–18

Habitat A small canal with soil banks occupied with *Phragmites australis* in the surroundings of the village near a group of blocks of flats and greenhouse agricultural lands. Pollution by artificial materials, mainly plastic, was observed.

Etymology Named after the mammalogist Nedko Nedyalkov (National Natural History Museum, Sofia, Bulgaria).

Description Shell pale corneous, glossy, transparent with fine growth lines (Fig. 5a,b). Comprises 3–4 whorls, which increase rapidly, regularly

with clearly visible to deep suture. Last whorl not deflected. First whorls immersed on both sides. Size small, 3.5–4.4 mm in diameter, 1.0–1.4 mm in height.

Animal Animal light grey with diffuse mantle pigmentation. Prostate gland bears 18–22 long diverticules (Fig. 5d). Phallotheca as long as praeputium (Fig. 5c). Bursa copulatrix cylindrical (Fig. 5d).

Remarks Due to the small shell (3.5–3.9 mm in diameter) (Fig. 6, Table 1) *Gyraulus nedyalkovi* n.



Figure 7 *Anisus leucostoma*, swamp east of Kazanli village: shell.

sp. resembles *G. piscinarum*, *G. ehrenbergi*, and *G. huwaizahensis*. The new species differs especially in the numbers of prostate diverticules, the main distinguishing feature between species in the genus *Gyraulus* (see: Meier-Brook, 1983), but also in shell characteristics.

Anisus leucostoma (Millet 1813)

Material examined Four specimens from a swamp east of Kazanli village ($N36^{\circ}48'24.7''$ $E34^{\circ}47'34.8''$, fig. 1.3).

A dissection revealed that the prostate gland bears 19 diverticules. This is in accord with Glöer & Meier-Brook (2008: 94) for *Anisus leucostoma*.

Remarks: Yıldırım et al. (2006) listed *Anisus spirorbis* from Eastern Anatolia and cited Boettger (1957) for reference. However, in their reference list they mentioned only O. Boettger (1905), overlooked C. R. Boettger (1957), who reported *Anisus spirorbis* from a swamp in Erzurum Province.

ACKNOWLEDGEMENT

We thank Nedko Nedyalkov (National Natural History Museum, Sofia, Bulgaria) who first took Dilian Georgiev on an expedition to Turkey on April 2009, and attracted his interest on investigations of the Turkish fauna. In addition we express our thanks to an anonymous reviewer how corrected the English and gave helpful comments which improved our paper.

REFERENCES

- ANGELOV AM 2000 *Mollusca: Gastropoda et Bivalvia aquae dulcis*. Catalogus Faunae Bulgaricae 4. 57 pp.
- BANK R 2004 Towards a catalogue and bibliography of the freshwater Mollusca of Greece. *Hedea* 6: 79–112.
- BOETTGER CR 1957 Über eine Ausbeute von Höhlenmollusken und einigen anderen Weichtieren aus der Türkei. *Archiv für Molluskenkunde* 86(1/3): 67–83.
- DAUTZENBERG P 1894 Liste des mollusques terrestres et fluviatiles recueillis par M. Th. Barrois en Palestine et en Syrie. *Revue biologique du Nord de la France* 6: 329–353.
- GLÖER P & BÖSSNECK U 2007 Zur Identität von *Gyraulus piscinarum* Bourguignat 1852 mit der Beschreibung von *G. bekaensis* n. sp. (Gastropoda: Planorbidae). *Mollusca* 25(2): 139–146.
- GLÖER P & FEHÉR Z 2004 *Bithynia leachii* (Sheppard, 1823) and *Bithynia troschelii* (Paasch, 1842) in Hungary (Prosobranchia: Bithyniidae). *Annales historico-naturales Musei nationalis Hungarici* 96: 285–297.
- GLÖER P & MEIER-BROOK C 2008 Redescription of *Anisus septemgyratus* (Rossmaessler, 1835) and *Anisus leucostoma* (Millet, 1813) (Gastropoda: Planorbidae). *Mollusca* 26(1): 89–94.
- GLÖER P & NASER MD 2007 *Gyraulus huwaizahensis* n. sp. – a new species from Mesopotamia, Iraq (Mollusca: Gastropoda: Planorbidae). *Mollusca* 25 (2): 147–152.
- GLÖER P & RÄHLE W 2009 *Gyraulus pamphylicus* n. sp. – a new species from Turkey (Mollusca: Gastropoda: Planorbidae). *Mollusca* 27(1): 57–60.
- GLÖER P & YILDIRIM MZ 2006 Some records of Bithyniidae from Turkey with the description of *Bithynia pesicci* n. sp. (Gastropoda: Bithyniidae). *Malakologische Abhandlungen* 24: 37–42.
- MEIER-BROOK C 1983 Taxonomic studies on *Gyraulus* (Gastropoda: Planorbidae). *Malacologia* 24 (1–2): 1–113.
- STURANY R 1904 Sitzung der mathematisch-naturwissenschaftlichen Klasse vom 21. April 1904. *Anzeiger der Kaiserlichen Akademie der Wissenschaften in Wien* 41: 115–118.
- YILDIRIM MZ 1999 Türkiye Prosobranchia (Gastropoda: Mollusca) Türleri ve Zoocoğrafik Yayınları. 1. Tatlı ve Aci Sular. *Turkish Journal of Zoology* 23 Ek Sayı 3: 877–900.
- YILDIRIM MZ, KOCA SB & KEBAPÇI Ü 2006 Supplement to the Prosobranchia (Mollusca: Gastropoda). Fauna of Fresh and Brackish Waters of Turkey. *Turkish Journal of Zoology* 30: 197–204.
- YILDIRIM MZ, GÜMÜŞ BA, KEBAPÇI Ü & KOCA SB 2006 The Basommatophoran Pulmonate Species (Mollusca: Gastropoda) of Turkey. *Turkish Journal of Zoology* 30: 445–458.
- YILDIRIM MZ & KEBAPÇI Ü 2006 Endemism of Land and Freshwater Gastropods in the Lakes Region (Turkey). *Muzeul Olteniei Craiova. Oltenia. Studii și comunicări. Științele Naturii* 25: 55–59.